# Abnormal Proactive and Reactive Cognitive Control During Conflict Processing in Major Depression

Marie-Anne Vanderhasselt, Rudi De Raedt, Annick De Paepe, and Kristien Aarts Ghent University Georges Otte and Jan Van Dorpe Ghent, Belgium

## Gilles Pourtois Ghent University

According to the Dual Mechanisms of Control framework, cognitive control consists of two complementary components: *proactive control* refers to anticipatory maintenance of goal-relevant information, whereas *reactive control* acts as a correction mechanism that is activated when a conflict occurs. Possibly, the well-known diminished inhibitory control in response to negative stimuli in Major Depressive Disorder (MDD) patients stems from a breakdown in proactive control, and/or anomalies in reactive cognitive control. In our study, MDD patients specifically showed increased response latencies when actively inhibiting a dominant response to a sad compared with a happy face. This condition was associated with a longer duration of a dominant ERP topography (800–900 ms poststimulus onset) and a stronger activity in the bilateral dorsal anterior cingulate cortex, reflecting abnormal reactive control when inhibiting attention to a negative stimulus. Moreover, MDD patients showed abnormalities in proactive cognitive control when preparing for the upcoming imperative stimulus (abnormal modulation of the contingent negative variation component), accompanied by more activity in brain regions belonging to the default mode network. All together, deficits to inhibit attention to negative information in MDD might originate from an abnormal use of both proactive resources and reactive control processes.

Keywords: Major Depressive Disorder, proactive control, reactive control, ERP topographic mapping analysis

Although Major Depressive Disorder (MDD) is primarily characterized by persistent low mood, recurrent negative thoughts and anhedonia, it is also accompanied by core cognitive deficits at the level of information processing. These impairments are most pronounced for negative information, with specific difficulties in inhibiting attention to task-irrelevant negative information (Joormann, Yoon, & Zetsche, 2007), leading to depressed mood. Depressive feelings have been associated with disregulated corticolimbic interactions accompanied by changes in dorsal neocortical activation, specifically the dorsolateral prefrontal cortex (DLPFC, Brodman Area (BA) 9/46), the dorsal anterior cingulate (dACC, BA 24), the inferior parietal cortex (IPC, BA 40) and the striatum (Mayberg, 1997). However, it remains unclear whether this dorsal activation is either decreased or increased during cognitive control operations engaged during the processing of emotional stimuli in currently or remitted depressed patients (Joormann & Gotfib, 2008). Whereas some studies reported hyperactivity in these dorsal regions in depressed patients during cognitive control operations (Chiu & Deldin, 2007; Harvey et al., 2005; Holmes & Pizzagalli, 2008a; Luu, Flaisch, & Tucker, 2000; Liotti, Woldorff, Perez, & Mayberg, 2000), other studies found depression-related hypoactivity in these same areas, including the DLPFC and the dACC (Holmes & Pizzagalli, 2008b; Vanderhasselt & De Raedt, 2009). This discrepancy might stem from the fact that these studies focused on different aspects of cognitive control, including proactive and reactive mechanisms (Braver, Gray, & Burgess, 2007).

Recent theoretical accounts (Braver, 2012) have proposed that cognitive control is not exclusively related to reactive mechanisms in response to an imperative stimulus (e.g., conflict), but that specific proactive processes may also play a role in conflict monitoring (and more generally goal-directed behavior). According to the Dual Mechanisms of Control framework (DMC) (Braver et al., 2007; Braver, 2012), proactive control refers to anticipatory or preparatory processes (i.e., activating and maintaining online goal-

Marie-Anne Vanderhasselt, Rudi De Raedt, Annick De Paepe, and Kristien Aarts, Department of Experimental Clinical and Health Psychology, Ghent University, Belgium; Georges Otte and Jan Van Dorpe, P.C. Dr. Guislain, Fr. Ferrerlaan 88A, 9000 Ghent – Belgium; Gilles Pourtois, Ghent University.

MAV (FW008/PD0/168) is a postdoctoral research fellow of the Research Foundation Flanders (FWO). Preparation of this paper was also supported by Grant BOF10/GOA/014 for a Concerted Research Action of Ghent University (awarded to RDR & GP). GP is supported by grants from the European Research Council (Starting Grant #200758) and Ghent University (BOF Grant #05Z01708). This work was supported by Grant P7/11 from the Interuniversity Attraction Poles program of the Belgian federal government.

Correspondence concerning this article should be addressed to Marie-Anne Vanderhasselt, Department of Experimental Clinical and Health Psychology, Ghent University, Henri Dunantlaan 2, 9000 Ghent, Belgium. E-mail: MarieAnne.Vanderhasselt@ugent.be

relevant information) aimed at enhancing coping with conflict before it actually takes place. This proactive control depends on contextual information (Miller & Cohen, 2001) and serves to guide the information processing system toward goal-relevant information before the onset of the imperative stimulus. On the other hand, reactive control refers to a correction mechanism that is activated when an ambiguous or conflict stimulus occurs (Jacoby, Kelley, & McElree, 1999). Reactive control mechanisms are essential to mobilize additional processing resources in order to eventually resolve this conflict. According to the DMC, although these two components of working memory and cognitive control likely operate at different moments during conflict monitoring (e.g., early/ sustained selection and late/transient correction for proactive and reactive mechanisms, respectively), they both depend on the integrity of a dorsal brain system comprising the DLPFC and dACC area.

Although most studies have focused on reactive modes of cognitive control in samples of depressed patients, some studies have also demonstrated abnormalities in proactive control associated with negative mood. An earlier study from West, Choi, and Travers (2010) observed that negative affect (measured using a Beck Depression Inventory) in healthy individuals was associated with attenuated proactive and reactive cognitive control (using a counting Stroop task). Moreover in clinically depressed patients, an abnormal contingent negative variation (CNV) during the engagement of preparatory processes has been observed (Ashton, Marshall, Hassanyeh, Marsh, & Wright-Honari, 1994; Heimberg et al., 1999; Giedke & Heimann, 1987). This slow cue-locked cortical potential, which is maximal over frontocentral sites, reflects anticipatory attention and effortful processing (Brunia & van Boxtel, 2001). Possibly, diminished inhibitory control in response to a (task-irrelevant) negative stimulus (as typically seen in depression) might stem from a breakdown in proactive control, which normally operates before the conflict is actually experienced (Braver, 2012). It appears therefore important to investigate both proactive and reactive modes of cognitive control in currently depressed patients.

In a recent Event Related Potential (ERP) study (Vanderhasselt et al., 2012), a new experimental paradigm to disentangle the respective contributions of these two cognitive control components during conflict monitoring was used: the Cued Emotional Conflict Task (CECT). Based on the presentation of a cue that informed participants about which S-R mapping to use later (either press, actual or opposite) when seeing a face stimulus using EEG methods, the amount of proactive control could be assessed during this time period preceding the onset of the imperative face stimulus (target). Derived from the DMC framework (Braver et al., 2007; 2012), proactive control was conceptualized as the active maintenance in working memory of a specific task goal. This is harder in the case of "opposite" and "actual" than "press," given that active emotion face discrimination is required in the former case, whereas only simple face detection is required in the latter case. Critically, the face had either a happy or sad expression, which made it possible to compare conflict processing of positive versus negative information, respectively.

The aim of the current study was to investigate proactive and reactive control mechanisms by comparing behavioral performance (reaction time (RT)) and electrophysiological effects (by means of ERP measurements) in MDD patients and healthy con-

trols during the CECT task. MDD patients were hypothesized to have selectively increased response latencies when encountering sad (compared with happy) faces that are preceded by the cue "opposite," because of a selective impairment in inhibiting a response to negative information in favor of the concurrent/alternative positive response. Using high-density ERP, we evaluated whether such a condition-specific behavioral deficit, if present in MDD, mainly results from selective abnormalities during proactive control (i.e., cue-related ERP activities), or instead during reactive control (i.e., ERP time-locked to the faces), compared with a group of matched healthy control participants. To address these questions, an advanced ERP topographic mapping analysis (Michel & Murray, 2012; Murray, Brunet, & Michel, 2008; Pourtois, Delplanque, Michel, & Vuilleumier, 2008) was used, combined with a standard distributed source localization method.<sup>1</sup> Using this data analysis, possible abnormalities in either proactive or reactive control during conflict monitoring in MDD can be delineated. Moreover, this study aims to gain insight into the brain networks involved in these processes and their potential alteration in MDD, with a specific focus on lateral and medial prefrontal cortex given the implication of these regions in proactive and reactive cognitive control effects.

#### Method

## **Protocol**

Participants who agreed to participate in the experiment were contacted by phone, and were screened on inclusion/exclusion criteria. Participants deemed to be eligible after this screening were invited for a clinical structural interview in the lab. Participants meeting all inclusion criteria were subsequently invited for an ERP session. Participants were asked not to smoke at least 2 hr before the start of the experiment. They gave their written informed consent and received 20 Euros for their participation. The study was approved by the medical ethics committee of the Ghent University hospital.

## **Participants**

Twenty individuals meeting the *DSM–IV* criteria for major depressive disorder (MDD) (13 females, mean age: 38.40, *SD*: 13.14), and 20 matched nondepressed (ND) individuals (15 females, mean age: 41.25, *SD*: 14.64), participated in this study.

The MDD ambulatory patients were recruited from a local Belgian psychiatric clinic and were diagnosed with MDD (four patients had a comorbidity with anxiety disorder; anxiety symptoms were secondary to a depressive illness). Before testing, the Hamilton Rating Scale for Depression (HAM-D; Hamilton, 1960) and the Mini-International Neuropsychiatric Interview (MINI; Sheehan et al., 1997), a structured clinical interview, were admin-

<sup>&</sup>lt;sup>1</sup> The added value of this data-driven clustering analysis is that it enables revealing condition-specific differences in the configuration of the ERP electric field (i.e., topography), which are difficult to capture otherwise using a standard peak analysis (i.e., latency or amplitude variations of specific ERP components evidenced at a few electrode positions), especially when substantial differences in the amplitude of the global ERP signal between groups (MDD vs. ND) can occur.

istered to examine the severity of the current MDD episode. The exclusion criteria for MDD patients were: (1) the presence of other mood disorders; (2) use of antipsychotics, monoamine oxidase inhibitors, tricyclic antidepressants and/or benzodiazepines; (3) a history of neurological conditions such as epilepsy, a brain trauma, loss of consciousness during more than 5 minutes; (4) a history of electroconvulsive therapy (ECT); (5) the abuse of alcohol during the last year; (6) a current or past substance dependence; (7) a current or past psychotic episode; and finally (8) the presence of learning disorders. Patients who had either serotonergic or noradrenargic antidepressants were included, but only if these patients received their medication for at least 2 weeks on a steady basis before testing. Healthy subjects were included in the control group for comparison purposes with the MDD patients if they were free of medication during the time of testing, and presented no evidence of current or past psychopathologic disorders (assessed using the MINI and HAM-D), or self-reported neurologic disorders or head injuries. This ND sample was matched with the depressed participants at the group level on sex, age and education.

All participants were native Caucasian Dutch speakers, had normal or corrected vision and were right-handed. Demographic and clinical characteristics of the participants are outlined in Table 1.

# Stimuli and Task

In the Cued Emotional Conflict Task (CECT) each trial starts with 1 out of 3 single written word cue presented in random order (see Figure 1; see Vanderhasselt et al., 2012 for details): "actual" to press a key corresponding to the emotional expression of the upcoming target face; "opposite" to respond to the opposite emotional expression of the target face; "press" to press a separate key when a face appeared, regardless of the emotional expression of the face (simple detection required). Fourteen faces (7F/7M) from the Karolinska Directed Emotional Faces data set (Lundqvist, Flykt, & Öhman, 1997) were used. Each of these faces was shown in a happy and a sad expression, in order to control for physical characteristics of the faces. Faces were followed by a blank screen that remained until a response was made. Participants were instructed to answer as quickly and as accurately as possible with one out of three fingers of their right dominant hand. Participants first completed 30 practice trials using five faces not shown during

Table 1Demographical and Clinical Data

the experimental blocks, followed by six blocks of 36 trials. Each block consisted of six trials of each cue/face combination, presented in random order.

## Questionnaires

Depressive symptoms were measured using the BDI-II (Beck, Steer, & Brown, 1967) and the HAM-D (Hamilton, 1960). The BDI-II is a 21-question, multiple-choice, self-report inventory, examining the severity and the occurrence of cognitive, affective, somatic and vegetative symptoms of depression during the last 2 weeks. The HAM-D is a semistructured interview, evaluating the severity of depression. The interview consists of 21 items and explores depressed mood, vegetative (e.g., insomnia, fatigue, anorexia) and cognitive symptoms and comorbid anxiety disturbances.

## **EEG Recording**

Continuous EEG was acquired using a 128-channel (pintype) Biosemi Active Two system (http://www.biosemi.com) referenced to the CMS-DRL ground with an analog bandpass. The data were digitized at a 24-bit resolution with a Least Significant Bit (LSB) value of 31.25 nV and a sampling rate of 512 Hz, using a low-pass fifth order sinc filter with a -3dB cutoff point at 102 Hz. ERPs of interest were computed offline following a standard sequence of data transformations (Picton et al., 2000): (1) -250/+1500 ms segmentation around the onset of word (cue) stimulus and -500/+2000 ms segmentation around the onset of the face (target) stimulus (2) prestimulus interval baseline correction (from -250 ms to the cue onset, and from -500 ms to target onset), (3) vertical ocular correction for blinks (Gratton, Coles, & Donchin, 1983) using the difference amplitude of two electrodes attached above and below the left eye, (4) artifact rejection [M = 84.47 trials]SEM = 1.98 amplitude scale ( $\mu V$ ) across participants: no difference between ND (M = 86.32 trials, SEM = 2.56) and MDD patients (M = 82.63 trials, SEM = 3.04) was evidenced; t = 0.93, p = .36], (5) averaging of trials, separately for each group (ND vs. MDD) and experimental condition (n = 6), and (6) 30 Hz low-pass digital filtering of the individual average data.

	ND (n = 19)		$\begin{array}{l} \text{MDD} \\ (n = 19) \end{array}$		Statistics
	Mean	SD	Mean	SD	t t
Age	40.11	14.09	37.89	13.30	0.50
% Female	70%	N/A	60%	N/A	0.68
Number of depressive episodes	N/A	N/A	2.63	1.30	N/A
Age of onset depression (in years)	N/A	N/A	32.79	12.62	N/A
Duration present episode (in months)	N/A	N/A	6.95	5.51	N/A
BDI-II	1.53	3.79	34.21	10.84	$-12.41^{*}$
HAM-D	0.21	0.54	28.26	5.04	$-24.11^{*}$

*Note.* BDI: Beck Depression Inventory-II; HAM-D: Hamilton Rating Scale for Depression. \* p < .01.



*Figure 1.* Schematic overview of the Cued Emotional Conflict Task (CECT). First, a cue is presented in the center of the screen ("actual," "opposite" or "press"), followed by a face with an emotional expression (either happy or sad), leading to six possible cue-face combinations/conditions. ISI = interstimulus interval; intertribal interval (ITI) = intertrial interval.

## **Topographical Analyses**

In order to capture ERP differences between ND and MDD patients, a detailed topographic mapping analysis of the ERP data was performed, following a conventional data-analysis scheme (Michel et al., 2001; Pourtois, Dan, Grandjean, Sander, & Vuilleumier, 2005a; Pourtois, Thut, de Peralta, Michel, & Vuilleumier, 2005b). To precisely characterize topographic modulations over time and across conditions, a standard spatial cluster analysis was used. This pattern analysis efficiently summarizes a complex ERP data set into a smaller number of dominant field configurations, previously referred to as functional microstates (Lehmann & Skrandies, 1979). The rationale and basic principles of this temporal segmentation method have been extensively described elsewhere (e.g., Murray, Brunet, & Michel, 2008). Following standard practice, a topographic pattern analysis was first performed on the grand-average ERP data from stimulus onset until 2000 ms after stimulus onset (1000 consecutive time frames at 512 Hz sampling rate), using a standard K-means cluster method (Pascual-Marqui, 2002). The optimal number of topographic maps explaining the whole data set was determined objectively using both cross-validation (Pascual-Marqui, 2002) and Krzanowski-Lai criteria (Tibshirani, Walther, & Hastie, 2001). The dominant scalp topographies (identified by the previous analysis) were then fitted back to the ERP data of each individual subject using spatial fitting procedures to quantitatively determine their representation across subjects and conditions. This procedure thus provides fine-grained quantitative values, such as the duration or strength (Global Field Power – GFP), which are critical estimates of the significance of a given topography, not available otherwise in a classical component analysis (Picton et al., 2000). The resulting duration or strength (GFP) values were entered in mixed ANOVAs with the between-subjects factor group (ND vs. MDD) and the within-subject factors emotion (happy, sad) and condition (actual, opposite, press). These analyses were carried out using CARTOOL software (Version 3.34; developed by D. Brunet, Functional Brain Mapping Laboratory, Geneva, Switzerland). Given that the cue locked CNV (as a function of the amount

of proactive control to be exerted; see below) has typically been associated with amplitude changes in previous ERP studies (see Brunia & van Boxtel, 2001), the CNV component was expected to vary in amplitude. On the other hand, no such prediction can be made regarding the expression of the reactive component (likely affecting the amplitude or latency of the ERP signal time-locked to the onset of the imperative face stimulus).

## **Source Localization Analyses**

To estimate the likely neural sources associated with the dominant electrical field configurations identified by the previous analyses, a specific distributed linear inverse solution was used, namely standardized low-resolution brain electromagnetic tomography (sLORETA, Pascual-Marqui, 2002).<sup>2</sup> A direct comparison between the inverse solution results for the opposite-sad and actual-sad condition (MDD

<sup>&</sup>lt;sup>2</sup> sLORETA is based on the neurophysiological assumption of coherent coactivation of neighboring cortical areas (known to have highly synchronized activity; see Silva, Amitai, & Connors, 1991) and, accordingly, it computes the "smoothest" of all possible activity distributions (i.e., no a priori assumption is made regarding the number and locations of the sources). Mathematical validation of this distributed source localization technique has been demonstrated (Sekihara, Sahani, & Nagarajan, 2005). sLORETA solutions are computed within a three-shell spherical head model coregistered to the MNI152 template (Mazziotta et al., 2001). The source locations were therefore given as (x, y, z) coordinates (x from left to right; y from posterior to anterior; z from inferior to superior). sLORETA estimates the three-dimensional intracerebral current density distribution in 6239 voxels (5 mm resolution), each voxel containing an equivalent current dipole. This three-dimensional solution space in which the inverse problem is solved, is restricted to the cortical gray matter (and hippocampus). The head model for the inverse solution uses the electric potential lead field computed with a boundary element method applied to the MNI152 template (Fuchs, Kastner, Wagner, Hawes, & Ebersole, 2002). Scalp electrode coordinates on the MNI brain are derived from the international 5% system (Jurcak, Tsuzuki, & Dan, 2007). The calculation of all reconstruction parameters was based on the computed common average reference. sLORETA units were scaled to ampere per square meter (A/m2).

patients) was performed separately for the cue and target-related activity, using paired *t* tests. To reveal significant effects, we used a stringent nonparametric randomization test (relying on 5000 iterations) using a corrected p < .05 value.

## Results

Because of technical problems, data from one healthy participant (n = 19) and one patient from the MDD group (n = 19) were omitted from the analyses of the behavioral and electrophysiological data.

## **Demographics and Self-Report Data**

Table 1 summarizes demographic, clinical and self-report data. Groups did not differ on any demographic variable (ps > .05).

## **Behavioral Data**

We refer to Figure 2 for an overview of median RT data for correct responses. Accuracy rates ranged between 90.69% and 96.88%, with no difference between groups, ts < 1.65 & ps > .1. Therefore, only trials for which participants made a correct response were included in the analysis. The *Cue* (opposite, actual, press) x *Emotion* (sad, happy) x *Group* (ND, MDD) ANOVA with the median RTs as dependent variable revealed a three-way interaction, F(2, 35) = 4.41, p = .02,  $\eta_p^2 = .20$  (also all the main effects were significant, Fs > 9.89, ps < .005, as well as the interaction between *Cue* x *Group*, F = 9.03, p < .001, and *Cue* x *Emotion*, F = 11.36, p < .001.

Follow-up independent *t* tests revealed that RT in MDD patients were not different from ND controls following the press trials, ts < 1.42, ps > .17. MDD patients demonstrated, on the other hand, increased response latencies on the four other CECT trials ("opposite-sad," "opposite-happy," "actual-sad" and "actualhappy"), ts > 4.18, ps < .001. However, as hypothesized, withingroup analysis revealed that MDD patients demonstrated greater RT to "opposite-sad" than "opposite-happy" trials, t(18) = 2.24, p = .04, whereas the ND participants had balanced RT for both CECT trials, t(18) = .78, p = .46. Both groups had greater RT on "actual-sad" than "actual-happy" trials, ps < .001, but did not differ in RT to "press-sad" and "press-happy" trials, ps > .31.

## **ERP** Data

Figure 3 presents, for each group separately, the grand average ERP waveforms at electrode FCZ for either cue/word or target/face-related activities.

## **Topographical Components**

**Target.** A spatiotemporal cluster analysis was performed on a large time-window (i.e., 2000 ms), encompassing the early (P1 and N170), midlatency (P2) and late (N2 and P3) ERP components generated in response to the happy or sad faces. A solution with 10 dominant maps/topographies explained 93% of the variance. Remarkably, consistent with our prediction, this cluster analysis disclosed a dominant topography that was diagnostic of the condition "opposite-sad," for the MDD group selectively. This field



*Figure 2.* Median RT data (and standard deviations) (in ms) for correct CECT trials, both in the ND (n = 19) and the MDD (n = 19) sample. MDD patients demonstrated significantly greater RT to "opposite-sad" than "opposite-happy" trials, whereas the ND participants had balanced RT for both CECT trials. Both groups had significantly greater RT on "actual-sad" than "actual-happy" trials but did not differ in RT to "press-sad" and "press-happy" trials.



*Figure 3.* Grand average ERP waveforms at electrode FCZ, separately for the cue/word (upper panels) and the target/face (lower panels). A, In the ND group, a clear modulation of the CNV component was evidenced at a late latency following cue onset, indicated by a larger amplitude for the "press" than both "actual" and "opposite." B, In the MDD group, no similar amplitude variation of the CNV component was visible. C, In the ND group, no systematic amplitude variation of the ERP signal was visible across the four main experimental conditions. D, By contrast, in the MDD group, starting 650 ms poststimulus onset, the opposite sad condition clearly elicited a differential ERP activity during a prolonged time interval (indicated by the vertical red arrows), compared with the three other experimental conditions.

configuration lasted  $\sim$ 400 ms, starting 650 ms following stimulus onset. This topography was characterized by a positive component over centro-parietal scalp leads, accompanied by a left-lateralized negative component over frontal electrode locations (see Figure 4A). Following standard practice, a fitting of this dominant map back to the individual ERP data was performed to verify, at the statistical level, whether this topography was diagnostic of the "opposite-sad" condition, selectively for the MDD group. The dependent variable was the degree of similarity between the grand average topographical template (identified by the previous analysis) and the single-subject data. The duration values, obtained for this dominant map after fitting, were submitted to a 2 (*Group*)  $\times$ 2 (*Emotion*)  $\times$  2 (*Cue*) mixed ANOVA. This analysis revealed a three-way interaction, F(2, 35) = 3.450, p = .04. Whereas for the ND group, the duration of this dominant topography did not vary depending on emotion and cue [interaction, F < .7, p > .54], it tended to do so in the MDD group [interaction, F(2, 17) = 2.96, p = .08]. As can be seen from Figure 4B, in the MDD group only, this topography had a longer duration for the condition "oppositesad" than for all other conditions. The direct comparisons between "opposite-sad" versus "actual sad," t(18) = 2.56, p = .02, and "opposite-sad" versus "opposite-happy," t(18) = 3.19, p < .01, confirmed a prolonged duration of this dominant topography selectively for the condition "opposite-sad."

Cue. A similar data-driven analysis was used to explore possible topographic changes across cues and between groups occurring before the onset of the imperative face stimuli, namely during the processing of the cue foreshadowing the type of visual categorization to be made by participants. The spatiotemporal cluster analysis was performed on a large time window (i.e., 1500 ms), encompassing the early, midlatency and late ERP components generated in response to the three possible cues ("press," "actual" or "opposite"). This analysis disclosed a solution with eight dominant topographies accounting for 93% of the variance. A visual inspection of these maps (ND group) showed that early sensory processing of the cue (bilateral P1 and N1 occipital components) was later followed, after a transition phase where sustained midlatency ERP components were generated, by a clear-cut CNV component (dominant topography), whose expression was most obvious starting 1000 ms following cue onset and showing a sustained effect until 1500 ms postcue onset. Therefore, this prolonged time interval (i.e., 1000-1500 ms postcue onset) was used to assess whether the dominant CNV topography (characterized by a fronto-central negative activity; see Figure 5A) underwent change in strength depending on the task demands/conditions, as well as a function of the group (ND vs. MDD). Interestingly, during this time interval, the amplitude of the CNV component appeared to be much reduced for the MDD patients, compared



Figure 4 (opposite)

with the control participants. Moreover, for the ND participants only, the amplitude of this CNV component varied according to the cue type, being the largest for the "press" instructions, but reduced alike for both the "opposite" and "actual" conditions. To corroborate these observations at the statistical level, the amplitude values, obtained for this dominant CNV map after fitting during this prolonged time interval postcue onset, were submitted to a 2  $(Group) \times 3$  (*Cue*) mixed ANOVA. This analysis showed mains effects of *Group*, F(1, 36) = 33.68, p < .001, and *Cue*, F(2, 35) =6.58, p < .01. Whereas the former effect confirmed that the strength of the CNV was substantially reduced for MDD patients compared with controls, the latter indicated a change in the amplitude of the CNV component depending on the cue type. Interestingly, as can be seen from Figure 5B, paired t tests showed that for the ND participants, the CNV had the largest amplitude for the press condition than either the "actual," t(18) = 2.80, p = .01, or "opposite" cue, t(18) = 2.31, p = .03. These two latter cues ("actual" and "opposite") were not different from one another, t(18) = 1.64, p = .12. In MDD patients, the CNV component was not different between cues "opposite" and "press," t(18) = 0.88, p = .39, suggesting that the preparatory processes (reflected by the CNV component) were similar in these two different conditions. Moreover, the CNV component in MDD patients was less negative in amplitude for "actual" compared with "press," t(18) = 2.28, p =.04, suggesting that their impairment during the cue period was not general (or generic), but mainly concerned the "opposite" cue condition (because no difference in amplitude between "press" and "opposite"). No difference for the amplitude of the CNV component was found between "actual" and "opposite" cue conditions, t(18) = 0.99, p = .34.

# **Inverse Solutions**

**Target.** A source localization analysis based on sLoreta showed that the configuration of the intracranial generators associated with this dominant topography (ERPs for the target faces) mainly involved bilateral dorsal medial frontal cortex brain areas, with a maximum found in Brodmann area 6 ( $X = \pm 5$ , Y = -25, Z = 54), with a notable spread of this broad activation toward more ventral medial frontal sites, including the ACC-Brodmann area 24 ( $X = \pm 4$ , Y = -17, Z = 45; see Figure 4C). Next, a direct statistical comparison was performed in the inverse solution space to establish whether this enhanced medial frontal cortex activation was significant for the "opposite-sad" condition or not. For this purpose, we compared for the MDD patients the process-

ing of the exact same face stimuli (namely sad faces) but when they were either associated with an opposite stimulus-response mapping ("opposite-sad") or the normal/intuitive one ("actualsad"). A 100 ms interval was selected during which the dominant topography (see above) associated with the "opposite-sad" condition in the MDD group was found to be maximal (i.e., 800–900 ms time-interval poststimulus onset). This contrast revealed a stronger bilateral dorsal ACC (BA 24) activation for "opposite-sad" than "actual-sad" ( $X = \pm 5$ , Y = -15, Z = 41; t(18) = 2.28, p < .025), stimuli.

**Cue.** During the time interval corresponding to the dominant CNV topography (1000-1500 ms postcue onset), the statistical comparison in the inverse solution space (sLoreta) was compared between the condition "opposite" versus "actual," for the MDD patients selectively. Because the CNV in MDD patients during the "opposite" condition (where proactive control was required) showed an abnormal response profile (e.g., it was not numerically different compared with the CNV recorded during the baseline control condition "press;" see above), this contrast enabled shedding light on possible impaired proactive or preparatory brain processes during the anticipation (foreperiod) for the "opposite" condition in the MDD group. This contrast revealed that "opposite" led to a larger activity than actual within a distributed network, involving mainly the right middle frontal gyrus (X = +21, Y = +22, Z = +44; t(18) = 2.41, p < .025], and the precuneus on both sides (X =  $\pm$  15, Y = -59, Z = +24; t(18) = 2.30, p < .025] (see Figure 5C). This contrast revealed a stronger bilateral dorsal ACC (BA 24) activation for "opposite-sad" than "actual-sad" (X =  $\pm 5$ , Y = -15, Z = 41; t(18) = 2.28, p<.025), stimuli.

### Discussion

The goal of the present study was to examine whether deficits to inhibit attention to negative information in MDD patients were related to anomalies in proactive and/or reactive cognitive control.

Behavioral results showed that, over all participants: 1) the processing of the actual emotion was faster for happy than sad faces; and 2) the difference between actual and opposite cues was greater for positive than negative faces. This pattern is consistent with a general positivity bias; that is, participants are faster at categorizing stimuli as positive than categorizing stimuli as negative. Importantly, in MDD only, behavioral data showed a selective slowing of responses when actively inhibiting a dominant response to a sad face and being required to use the alternative

*Figure 4 (opposite).* A, Grand average ERP data (opposite sad, MDD patients) time-locked to the onset of the imperative (sad) face stimulus and shown using a standard butterfly plot (overlaid traces), including all 128 channels. The vertical dashed line indicates the onset of the imperative face stimulus. The topographical segmentation analysis showed that a dominant and diagnostic scalp configuration had a prolonged duration ~650–1050 ms postface onset (highlighted by the shaded orange frame), in this condition, selectively for the MDD patients. This topography was characterized by a dipolar field including a left-lateralized lateral (pre)fontral negativity and a central posterior-parietal positivity. B, The fitting of this dominant topography (see Method) revealed a condition- and group-specific effect. The duration of this topography was the longest in the opposite sad condition than the other experimental conditions in the MDD group, with no such variation seen across conditions for the ND group. \* p < .05; \*\* p < .01. C, Source localization results for this dominant topography. A direct statistical comparison (MDD patients) in the inverse solution space (sLoreta) between opposite sad and actual sad (800–900 ms postface stimulus onset) revealed a stronger dorsal ACC activity in the former than the latter condition. A sagital and coronal views are provided.



Figure 5 (opposite)

(opposite) response mapping, that is, to press the happy face response key. This emotion-specific deficit in MDD is in accordance with prior research (e.g., Joormann & Gotfib, 2008), and indicates that these patients encounter selective difficulties in overriding habitual (dominant) responses to negative information and in turn select an alternative (and counterintuitive) stimulusresponse mapping in this condition.

For this "opposite-sad" condition, analyses of the ERP data time-locked to the onset of the target faces - revealed a significant longer duration of a dominant topography than the other conditions in MDD patients. The reconstructed intracranial generators of this dominant topography involved bilateral dorsal medial frontal areas, with a spread toward ventral medial frontal sides, including the ACC. Crucially, this ACC area was significantly more active (800-900 ms poststimulus onset) for MDD patients when they were required to process sad faces but had to categorize them as positive (happy), the condition eliciting the maximum interference (and hence conflict) at the behavioral level in these patients. The ACC is known to be a critical hub for performance monitoring, and - following dominant models - conflict-related ACC activations usually reflect the need to exert additional top-down control in the face of conflict or error (Botvinick, Braver, Barch, Carter, & Cohen, 2001). Possibly, the current ERP results suggest that MDD patients needed to spark enhanced conflict-related dorsal ACC activity following the onset of sad faces that had to be categorized as happy faces, in order to overcome this strong interference effect. Although the reason for the increased ACC activity in this "opposite-sad" condition is unclear at the moment (i.e., it could reflect either compensatory processes or enhanced conflict detection), this result confirms the assumption of abnormal reactive control in major depression (Drevets, Price, & Furey, 2008), specifically when inhibiting attention to negative information.

Noteworthy, relative to ND participants, our results also showed proactive control abnormalities in MDD patients. In the ND group, the CNV component was found largest (i.e., more negative) when these healthy individuals anticipated a simple detection task ("press"), than the two other conditions requiring active maintenance of a complex task goal in working memory and hence, enhanced proactive control ("actual" and "opposite"). This association between CNV amplitudes and task demands or working memory load expressed by the cue is consistent with earlier ERP results in the literature (McEvoy, Smith, & Gevins, 1998; Gevins et al., 1996; Tecce, 1972). Interestingly, the current ERP results showed that the cues "actual" and "opposite" had a comparable CNV amplitude change, which is in line with the DMC framework. This account states that "Under proactive control conditions, prefrontal cortex activity should be present reliably across events, and not just on those in which it is most needed" (Braver et al., 2007, pp. 89). In sum, these new electrophysiological findings suggest that healthy controls used, based on the specific instructional cue, a proactive strategy to actively maintain goalrelevant information in working memory.

Strikingly, no such amplitude modulation of the CNV component (as was observed in the ND) was evidenced for MDD patients (i.e., only the cue "actual" led to a lower CNV amplitude than the cue "press," whereas the cue "opposite" was associated with a CNV that was equally large as for the cue "press"). Even though research is scarce on this topic, prior studies reported abnormal CNV amplitudes in individuals with MDD (Ashton et al., 1994; Ashton et al., 1988; Giedke & Bolz, 1980; Timsit-Berthier, 1993). The reasons as to why abnormal proactive control is "spontaneously" exerted by MDD patients during the foreperiod are not entirely clear yet at this stage. However, the possibility that this effect is simply caused by an overall breakdown in motivation or a cease in cognitive processing in MDD can be ruled out. This is because MDD patients showed a condition-specific reactive effect, that is, a differentiation between "opposite-sad" and "oppositehappy" trials at the behavioral and electrophysiological levels. Interestingly in the current study, CNV amplitudes in MDD patients were robustly decreased in general (i.e., more positive amplitudes for the three cue types). Such decreased CNV amplitudes have been related to problematic attentional resource allocation caused by mind wandering, daydreaming or active distraction by some other task (Tecce, 1972; Travis & Tecce, 1998; Tecce & Cattanach, 1993; Travis, Tecce, Arenander, & Wallace, 2002; Travis, Tecce, & Guttman, 2000). Moreover, it is also assumed that CNV amplitudes reflect deficient ability to engage attention to the current task goal, and away from potentially depressogenic thinking or rumination (Bostanov et al., 2012). These assumptions are in line with our current research findings, for which - in the MDD only - the cue "opposite" led to a significantly larger activity than cue "actual" within a distributed network, involving mainly the right middle frontal gyrus and the precuneus on both sides. Interestingly, these two nonoverlapping brain regions are typically related to the default mode network, being mostly responsive at rest and hypothesized to generate spontaneous thoughts or internally guided (as opposed to external-driven) mental processes, such as mind wandering (Raichle et al., 2001). Possibly, engaging in internal guided mental processing, such as mind wandering, might have distracted MDD patients from anticipating the upcom-

*Figure 5 (opposite).* A, Grand average ERP data (opposite sad, MDD patients) time-locked to the onset of the imperative (sad) face stimulus and shown using a standard butterfly plot (overlaid traces), including all 128 channels. The vertical dashed line indicates the onset of the imperative face stimulus. The topographical segmentation analysis showed that a dominant and diagnostic scalp configuration had a prolonged duration ~650–1050 ms postface onset (highlighted by the shaded orange frame), in this condition, selectively for the MDD patients. This topography was characterized by a dipolar field including a left-lateralized lateral (pre)fontral negativity and a central posterior-parietal positivity. B, The fitting of this dominant topography (see Method) revealed a condition- and group-specific effect. The duration of this topography was the longest in the opposite sad condition than the other experimental conditions in the MDD group, with no such variation seen across conditions for the ND group. \* p < .05; \*\* p < .01. C, Source localization results for this dominant topography. A direct statistical comparison (MDD patients) in the inverse solution space (sLoreta) between opposite sad and actual sad (800–900 ms postface stimulus onset) revealed a stronger dorsal ACC activity in the former than the latter condition. A sagital and coronal views are provided.

ing imperative stimulus, leading to inefficient proactive control. Presumably, an exaggerated internal focus, which is known to consume important processing resources (Jones, Siegle, Muelly, Haggerty, & Ghinassi, 2010), prevented a sharpening of their cognitive resources during the foreperiod, although these patients never stopped processing during the anticipation period. Alternatively, as it is assumed that a more negative CNV amplitude corresponds to more cognitive efforts (McEvoy, Smith, & Gevins, 1998; Gevins et al., 1996; Tecce, 1972), this CNV pattern might reflect the fact that MDD patients, compared with ND, were perhaps exerting too much proactive control. Hence, in this scenario, MDD patients would show abnormal cognitive control because of an exaggerated or too high level of proactive control during conflict anticipation. In sum, these cue-locked findings suggest abnormal anticipation in MDD, that is, abnormalities to maintain online, into short-term memory, active goal relevant representations to adjust cognitive resources based on the specific instructional cue. Future studies are needed to establish whether this effect corresponds either to an abnormally low or high proactive control in these patients.

All together, our results provide evidence that the observed deficit to inhibit attention to negative information in MDD found at the behavioral level, might stem from a combination of abnormal proactive and reactive control mechanisms. The DMC framework posits that both modes of control are complementing, and the computational tradeoff between both depends on individual differences. More specifically, based on this model, individuals scoring high on the Behavioral Approach System (BAS) show an increased tendency to use proactive cognitive control, whereas individuals scoring high on Behavioral Inhibition System (BIS) show an increased predisposition to use reactive control. It is well known that MDD patients are characterized by lowered BAS but overactive BIS activations (Kasch, Rottenberg, Arnow, & Gotlib, 2002). In line with predictions of the DMC account, the present findings suggest that MDD patients are less able than nondepressed individuals to proactively use effective cognitive control resources during the foreperiod foreshadowing the onset of the imperative face stimulus. The DMC further suggests that affectrelated traits influence the cost-benefit balance between proactive versus reactive control during goal-directed performance monitoring (Braver, 2012). Although the present findings do not offer statistical evidence for a tradeoff between both modes of cognitive control in MDD in the "opposite-sad" condition (r19 = .14, p =.55 for the correlation between CNV topography and LPC topography), it might be that - based on predictions from the DMC inefficient proactive control leads to an increased need of reactive control in order to overcome a strong interference at the behavioral level. In other words, MDD patients might be less able than nondepressed individuals to proactively employ cognitive control resources during the foreperiod foreshadowing the onset of the imperative face stimulus, which would result in a greater conflict in case this stimulus carries an interfering negative emotional expression. The nonsignificant correlation between CNV and LPC (as observed here) does not contradict this latter prediction derived from the DMC framework, however. Indeed, neither the CNV nor the LPC is uniquely related to cognitive control processes (proactive and reactive, respectively). Instead, each of these two ERP components/topographies appears to capture some variance related to cognitive control (besides other processes), but does not equate

it. Accordingly, finding a significant correlation between these two distant neural events during cognitive control and conflict monitoring appears unlikely given that these two ERP components are not reflecting pure measures of these cognitive processes. Further research is needed to establish whether abnormal reactive control effects regarding the inhibition of attention to negative information in depression could be predicted by systematic and traceable neurophysiological changes taking place earlier in time and corresponding to proactive control processes, as put forward in the DMC framework.

Some limitations of the present study should be mentioned. The majority of the MDD patients were on antidepressant medication (+/-80%), which might have influenced cognitive functioning, even though medication alone cannot a priori account for interaction effects between cue type and emotional face content reported in our study. Moreover, only patients who had either serotonergic or noradrenargic antidepressants were included, for at least 2 weeks before testing, hence ruling out the possibility of acute effects of these specific drugs on the reported behavioral and/or electrophysiological results. Also, patients who had received antipsychotics, monoamine oxidase inhibitors, tricyclic antidepressants and/or benzodiazepines, medications which are known to be associated with cognitive impairments, were excluded. Finally, although we are confident that these behavioral and ERP results cannot be explained easily by task difficulty alone, it remains challenging to demonstrate modulatory effects of depression on cognitive control and conflict detection, which are eventually fully orthogonal to changes in task difficulty. Given that conflict typically arises in situations where interference is created and for which RT and error rate will by definition increase, it is important to ascertain that depression influences conflict detection processes specifically, rather than taxing difficult trials more generally.

In sum, our results showed that the observed inhibitory deficit for negative information in MDD found at the behavioral level might stem from inefficient proactive control (possibly because of excessive internally focused processing that consumes important processing resources) and an abnormal reactive control.

## References

- Ashton, C. H., Marshall, E. F., Hassanyeh, F., Marsh, V. R., & Wright-Honari, S. (1994). Biological correlates of deliberate self-harm behaviour: A study of electroencephalographic, biochemical and psychological variables in parasuicide. *Acta Psychiatrica Scandinavica*, 90, 316–323. doi: 10.1111/j.1600-0447.1994.tb01600.x
- Ashton, H., Golding, J. F., March, V. R., Thompson, J. W., Hassanyeh, F., & Tyrer, S. P. (1988). Cortical evoked potentials and clinical rating scales as measures of depressive illness. *Psychological Medicine*, 18, 305–317. doi:10.1017/S0033291700007856
- Beck, A. T., Steer, R. A., & Brown, G. K. (1967). Manual for the Beck Depression Inventory-II. San Antonio, TX: Psychological Corporation.
- Bostanov, V., Keune, P. M., Kotchoubey, B., & Hautzinger, M. (2012). Event-related brain potentials reflect increased concentration ability after mindfulness-based cognitive therapy for depression: A randomized clinical trial. *Psychiatry Research*, 199, 174–180.
- Botvinick, M. M., Braver, T. S., Barch, D. M., Carter, C. S., & Cohen, J. D. (2001). Conflict monitoring and cognitive control. *Psychological Review*, 108, 624–652. doi:10.1037/0033-295X.108.3.624
- Braver, T. S. (2012). The variable nature of cognitive control: A dual mechanisms framework. *Trends in Cognitive Sciences*, 16, 106–113. doi:10.1016/j.tics.2011.12.010

- Braver, T. S., Gray, J. R., & Burgess, G. C. (2007). Explaining the many varieties of working memory variation: Dual mechanisms of cognitive control. In A. R. A. Conway, C. Jarrold, M. J. Kane, A. Miyake, & J. N. Towse (Eds.), *Variation in working memory* (pp. 76–106). Oxford, England: Oxford University Press.
- Brunia, C. H. M., & van Boxtel, G. J. M. (2001). Wait and see. *Interna*tional Journal of Psychophysiology, 43, 59–75. doi:10.1016/S0167-8760(01)00179-9
- Chiu, P. H., & Deldin, P. J. (2007). Neural evidence for enhanced error detection in major depressive disorder. *American Journal of Psychiatry*, 164, 608–616. doi:10.1176/appi.ajp.164.4.608
- Drevets, W. C., Price, J. L., & Furey, M. L. (2008). Brain structural and functional abnormalities in mood disorders: Implications for neurocircuitry models of depression. *Brain Structure Function*, 213, 93–118. doi:10.1007/s00429-008-0189-x
- Fuchs, M., Kastner, J., Wagner, M., Hawes, S., & Ebersole, J. S. (2002). A standardized boundary element method volume conductor model. *Clinical Neurophysiology*, 113, 702–712. doi:10.1016/S1388-2457(02)00030-5
- Gevins, A., Smith, M. E., Le, J., Leong, H., Bennett, J., Martin, N., . . . Whitfield, S. (1996). High resolution evoked potential imaging of the cortical dynamics of human working memory. *Electroencephalogram Clinical Neurophysiology*, 98(4), 327–348. doi:10.1016/0013-4694(96)00288-X
- Giedke, H., & Bolz, J. (1980). Pre- and postimperative negative variation (CNV and PINV) under different conditions of controllability in depressed patients and healthy controls. *Progress in Brain Research*, 54, 579–582. doi:10.1016/S0079-6123(08)61679-9
- Giedke, H., & Heimann, H. (1987). Psychophysiological aspects of depressive syndromes. *Pharmacopsychiatry*, 20, 177–180. doi:10.1055/s-2007-1017098
- Gratton, G., Coles, M. G. H., & Donchin, E. (1983). A new method for off-line removal of ocular artefact. *Electroencephalography and Clinical Neurophysiology*, 55, 468–484. doi:10.1016/0013-4694(83)90135-9
- Hamilton, M. (1960). A rating scale for depression. Journal of Neurology Neurosurgery and Psychiatry, 23, 56–62. doi:10.1136/jnnp.23.1.56
- Harvey, P. O., Fossati, P., Pochon, J. B., Levy, R., LeBastard, G., Leheriey, S., & Dubois, B. (2005). Cognitive control and brain resources in major depression: An fMRI study using the n-back task. *NeuroImage*, 26(3), 860–869. doi:10.1016/j.neuroimage.2005.02.048
- Heimberg, D. R., Naber, G., Hemmeter, U., Zechner, S., Witzke, W., Gerhard, U., . . . Hobi, V. (1999). Contingent negative variation and attention in schizophrenic and depressed patients. *Neuropsychobiology*, 39, 131–140. doi:10.1159/000026578
- Holmes, A. J., & Pizzagalli, D. A. (2008a). Response conflict and frontocingulate dysfunction in unmedicated participants with major depression. *Neuropsychologia*, 46(12), 2904–2913. doi:10.1016/j.neuropsychologia.2008.05.028
- Holmes, A. J., & Pizzagalli, D. A. (2008b). Spatiotemporal dynamics of error processing dysfunctions in major depressive disorder. *Archives of General Psychiatry*, 65(2), 179–188. doi:10.1001/archgenpsychiatry .2007.19
- Jacoby, L. L., Kelley, C. M., & McElree, B. D. (1999). The role of cognitive control: Early selection versus late correction. In S. Chaiken & E. Trope (Eds.), *Dual process theories in social psychology* (pp. 383– 400). New York: Guilford Press.
- Jones, N. P., Siegle, G. J., Muelly, E. R., Haggerty, A., & Ghinassi, F. (2010). Poor performance on cognitive tasks in depression: Doing too much or not enough? *Cognitive Affective Behavioral Neuroscience*, 10, 129–140. doi:10.3758/CABN.10.1.129
- Joormann, J., & Gotfib, I. H. (2008). Updating the contents of working memory in depression: Interference from irrelevant negative material. *Journal of Abnormal Psychology*, 117, 182–192. doi:10.1037/0021-843X.117.1.182

- Joormann, J., Yoon, K. L., & Zetsche, U. (2007). Cognitive inhibition in depression. *Applied Preventive Psychology*, 12, 128–139. doi:10.1016/ j.appsy.2007.09.002
- Jurcak, V., Tsuzuki, D., & Dan, I. (2007). 10/20, 10/10, and 10/5 systems revisited: Their validity as relative head-surface-based positioning systems. *NeuroImage*, 34(4), 1600–1611. doi:10.1016/j.neuroimage.2006 .09.024
- Kasch, K. L., Rottenberg, J., Arnow, B. A., & Gotlib, I. H. (2002). Behavioral activation and inhibition systems and the severity and course of depression. *Journal of Abnormal Psychology*, *111*, 589–597. doi: 10.1037/0021-843X.111.4.589
- Liotti, M., Woldorff, M. G., Perez, R., & Mayberg, H. S. (2000). An ERP study of the temporal course of the Stroop color-word interference effect. *Neuropsychologia*, 38(5), 701–711. doi:10.1016/S0028-3932(99)00106-2
- Lundqvist, D., Flykt, A., & Öhman, A. (1997). *The Karolinska Directed Emotional Faces (KDEF)*. Department of Neurosciences, Karolinska Hospital, Stockholm.
- Luu, P., Flaisch, T., & Tucker, D. M. (2000). Medial frontal cortex in action monitoring. *Journal of Neuroscience*, 20(1), 464–469.
- Mayberg, H. S. (1997). Limbic-cortical dysregulation: A proposed model of depression. *Journal of Neuropsychiatry and Clinical Neurosciences*, 9(3), 471–481.
- Mazziotta, J., Toga, A., Evans, A., Fox, P., Lancaster, J., Zilles, K., . . . Mazoyer, B. (2001). A probabilistic atlas and reference system for the human brain: International Consortium for Brain Mapping (ICBM). *Philosophical Transactions of the Royal Society B-Biological Sciences*, 356.
- McEvoy, L. K., Smith, M. E., & Gevins, A. (1998). Dynamic cortical networks of verbal and spatial working memory: Effects of memory load and task practice. *Cerebral Cortex*, 8, 563–574. doi:10.1093/cercor/8.7 .563
- Michel, C. M., & Murray, M. M. (2012). Towards the utilization of EEG as a brain imaging tool. *NeuroImage*, 61, 371–385. doi:10.1016/j .neuroimage.2011.12.039
- Michel, C. M., Thut, G., Morand, S., Khateb, A., Pegna, A. J., de Peralta, R. G., . . . Landis, T. (2001). Electric source imaging of human brain functions. *Brain Research Reviews*, 36, 108–118. doi:10.1016/S0165-0173(01)00086-8
- Miller, E. K., & Cohen, J. D. (2001). An integrative theory of prefrontal cortex function. *Annual Review of Neuroscience*, 24, 167–202. doi: 10.1146/annurev.neuro.24.1.167
- Murray, M. M., Brunet, D., & Michel, C. M. (2008). Topographic ERP analyses: A step-by-step tutorial review. *Brain Topography*, 20, 249– 264. doi:10.1007/s10548-008-0054-5
- Pascual-Marqui, R. D. (2002). Standardized low-resolution brain electromagnetic tomography (sLORETA) technical details. *Methods and Findings in Experimental and Clinical Pharmacology*, 24, 5–12.
- Picton, T. W., Bentin, S., Berg, P., Donchin, E., Hillyard, S. A., Johnson, R., . . Taylor, M. J. (2000). Guidelines for using human event-related potentials to study cognition: Recording standards and publication criteria. *Psychophysiology*, 37, 127–152. doi:10.1111/1469-8986.3720127
- Pourtois, G., Dan, E. S., Grandjean, D., Sander, D., & Vuilleumier, P. (2005a). Enhanced extrastriate visual response to bandpass spatial frequency filtered fearful faces: Time course and topographic evokedpotentials mapping. *Human Brain Mapping*, 26, 65–79. doi:10.1002/ hbm.20130
- Pourtois, G., Delplanque, S., Michel, C., & Vuilleumier, P. (2008). Beyond conventional event-related brain potential (ERP): Exploring the time-course of visual emotion processing using topographic and principal component analyses. *Brain Topography*, 20, 265–277. doi: 10.1007/s10548-008-0053-6
- Pourtois, G., Thut, G., de Peralta, R. G., Michel, C., & Vuilleumier, P. (2005b). Two electrophysiological stages of spatial orienting towards

fearful faces: Early temporo-parietal activation preceding gain control in extrastriate visual cortex. *NeuroImage*, *26*, 149–163. doi:10.1016/j .neuroimage.2005.01.015

- Raichle, M. E., MacLeod, A. M., Snyder, A. Z., Powers, W. J., Gusnard, D. A., & Shulman, G. L. (2001). A default mode of brain function. *Proceedings of the National Academy of Sciences, USA of the United States of America*, 98, 676–682. doi:10.1073/pnas.98.2.676
- Sekihara, K., Sahani, M., & Nagarajan, S. S. (2005). Localization bias and spatial resolution of adaptive and non-adaptive spatial filters for MEG source reconstruction. *NeuroImage*, 25(4), 1056–1067. doi:10.1016/j .neuroimage.2004.11.051
- Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Janavs, J., Weiller, E., Keskiner, A., . . . Dunbar, G. C. (1997). The validity of the Mini International Neuropsychiatric Interview (MINI) according to the SCID-P and its reliability. *European Psychiatry*, 12, 232–241. doi: 10.1016/S0924-9338(97)83297-X
- Silva, L. R., Amitai, Y., & Connors, B. W. (1991). Intrinsic oscillations of neocortex generated by layes-5 pyramid neurons. *Science*, 251, 432– 435. doi:10.1126/science.1824881
- Tecce, J. J. (1972). Contingent negative variation (CNV) and psychological processes in man. *Psychological Bulletin*, 77, 73–108. doi:10.1037/ h0032177
- Tecce, J. J., & Cattanach, L. (1993). Contingent negative variation (CNV). In E. Niedermeyer & F. Lopes da Silva (Eds.), *Electroencephalography: Basic principles clinical applications and related fields*, 3rd ed. (pp. 887–910). Baltimore: Williams and Wilkins.
- Tibshirani, R., Walther, G., & Hastie, T. (2001). Estimating the number of clusters in a data set via the gap statistic. *Journal of the Royal Statistical Society Series B-Statistical Methodology*, 63, 411–423. doi:10.1111/ 1467-9868.00293

Timsit-Berthier, M. (1993). Contingent negative variation and its relation-

ships to arousal and stress in psychopathology. In W. C. McCallum & S. H. Curry (Eds.), *Slow Potential Changes in the Human Brain*. Plenum Press, pp. 219–227. doi:10.1007/978-1-4899-1597-9\_14

- Travis, F., & Tecce, J. J. (1998). Effects of distracting stimuli on CNV amplitude and reaction time. *International Journal of Psychophysiology*, 31, 45–50. doi:10.1016/S0167-8760(98)00037-3
- Travis, F., Tecce, J., Arenander, A., & Wallace, R. K. (2002). Patterns of EEG coherence, power, and contingent negative variation characterize the integration of transcendental and waking states. *Biological Psychol*ogy, 61, 293–319. doi:10.1016/S0301-0511(02)00048-0
- Travis, F., Tecce, J. J., & Guttman, J. (2000). Cortical plasticity, contingent negative variation, and transcendent experiences during practice of the Transcendental Meditation technique. *Biological Psychology*, 55, 41–55. doi:10.1016/S0301-0511(00)00063-6
- Vanderhasselt, M. A., & De Raedt, R. (2009). Impairments in cognitive control persist during remission from depression and are related to the number of past episodes: An event related potentials study. *Biological Psychology*, 81(3), 169–176. doi:10.1016/j.biopsycho.2009.03.009
- Vanderhasselt, M. A., De Raedt, R., Dillon, D. G., Dutra, S. J., Brooks, N., & Pizzagalli, D. A. (2012). Decreased cognitive control in response to negative information in patients with remitted depression: An eventrelated potential study. *Journal of Psychiatry and Neuroscience*, 37, 250–258. doi:10.1503/jpn.110089
- West, R., Choi, P., & Travers, S. (2010). The influence of negative affect on the neural correlates of proactive and reactive cognitive control. *International Journal of Psychophysiology*, 76, 107–117. doi:10.1016/j .ijpsycho.2010.03.002

Received April 16, 2013

Revision received November 14, 2013

Accepted November 18, 2013 ■